



TITLE:

ON THE ASYMPTOTIC BEHAVIOR OF
SOLUTIONS FOR THE DISCRETE
BOLTZMANN EQUATION WITH LINEAR AND
QUADRATIC TERMS(Mathematical Analysis
of Phenomena in Fluid and Plasma
Dynamics)

AUTHOR(S):

YAMAZAKI, MITSURU

CITATION:

YAMAZAKI, MITSURU. ON THE ASYMPTOTIC BEHAVIOR OF SOLUTIONS FOR THE DISCRETE BOLTZMANN EQUATION WITH LINEAR AND QUADRATIC TERMS(Mathematical Analysis of Phenomena in Fluid and Plasma Dynamics). 数理解析研究所講究録 1994, 862: 46-55

ISSUE DATE:

1994-03

URL:

<http://hdl.handle.net/2433/83866>

RIGHT:

ON THE ASYMPTOTIC BEHAVIOR OF SOLUTIONS FOR THE DISCRETE BOLTZMANN EQUATION WITH LINEAR AND QUADRATIC TERMS

MITSURU YAMAZAKI (山崎 満)

Dept. Mathematical Science, Univ. Tokyo

In this paper, we study the discrete Boltzmann equation in one-dimensional space with linear and quadratic terms. This system, which is different from the usual one by the intervention of linear terms, describes the gas motion of molecules which take only a finite number of velocities under the interactions between particles represented by the quadratic terms and also under the reflection of molecules at the inner wall of an infinite thin tube, represented by the linear terms which we treated in the papers [13], [14], [15], [16], [17], [18].

$$(B) \quad \begin{cases} \frac{\partial u_i}{\partial t} + c_i \frac{\partial u_i}{\partial x} = Q_i(u) + L_i(u) , \\ u_i|_{t=0} = u_i^0(x) . \end{cases}$$

The physical theory imposes to this system the natural conditions :

Conditions .—

$$(*) \quad \begin{cases} A_{ij}^{k\ell} \geq 0, & A_{ij}^{k\ell} = A_{ji}^{k\ell} = A_{ij}^{\ell k} , \\ A_{k\ell}^{ij} \neq 0 & \Rightarrow i \neq j , \\ \alpha_i^k \geq 0 & \text{and } \alpha_i^i = 0 \quad \text{for all } i, k . \end{cases}$$

This linear terms are more general than the ones which are obtained by considering solutions around constant stationary solutions, called constant Maxwellian. We suppose furthermore the conservation of momentum in the course of interactions and also reflections

Condition mvQ.—

$$A_{ij}^{k\ell} \neq 0 \quad \Rightarrow \quad c_i + c_j = c_k + c_\ell .$$

and

Condition mvL.—

$$\forall i \in I, \sum_{k \in I} \alpha_k^i (c_k - c_i) = 0.$$

Remarque : We omit here all detailed explanations of these conditions which are described in [18].

Remarque : We know already that for bounded, summable and positive Cauchy data, the solutions exist globally in time, they are positive and bounded on $[0, \infty)$ [13], [16], [18].

Under these assumptions, we show ‘asymptotic’ behaviors of solutions which means that, for bounded and summable Cauchy data, the solutions $\sum_{c_a=c_i} u_i(x+c_i t, t)$ tend almost everywhere and in $L^q(q \in [1, \infty))$ to a function $\varphi_a(x)$. Furthermore, supposing that the velocities are mutually different, we prove that, for i belonging to a subset I_0 , the $u_i(x+c_i t, t)$ converge in L^∞ to $\varphi_i(x) \equiv 0$. Finally we treat the small data case, supposing that “sufficiently” reflection coefficients α_i^k are non zero, which is incompatible with the momentum conservation for the linear terms (mvL). Then we show, according to the argument due to Shizuta and Kawashima [9], [11], the decay in $(1+t)^{-\frac{1}{4}}$ of solutions. At the end, we prove that this assumption is also necessary for the decay of solutions.

Definition.— We define a subset I_0 as follows :

$$(1) \quad I_0 = \{i : \exists j \text{ such that } \alpha_j^i > 0\}.$$

We denote $\gamma_0 = \max_{i \in I} c_i$ and $E_0 = \{i : c_i = \gamma_0\}$, then $\gamma_1 = \max_{i \in I \setminus E_0} c_i$ and $E_1 = \{i : c_i = \gamma_1\}$, \dots . In this way, we have the decreasing sequence of velocities $\gamma_0 > \gamma_1 > \dots$ and $E_a = \{i : c_i = \gamma_a\}$. We put then

$$(2) \quad U_a(x, t) = \sum_{c_i=\gamma_a} u_i(x, t)$$

and

$$(3) \quad U_a^0(x, t) = \sum_{c_i=\gamma_a} u_i^0(x, t) \quad \text{and} \quad \mu = \sum_{i \in I} \int_{\mathbf{R}} u_i^0(x) dx.$$

Lemma 1.— Suppose the conditions (mvQ) and (mvL). Let u_i^0 be positive, summable and bounded Cauchy data. Then we have

$$(4) \quad U_a(x + \gamma_a t, t) = U_a^0(x) + \int_0^t p_a(x + \gamma_a s, s) ds - \int_0^t n_a(x + \gamma_a s, s) ds$$

with $p_a(x, t) \geq 0$, $n_a(x, t) \geq 0$ and $\int_0^\infty \int_{\mathbf{R}} p_a(x, t) dx dt \leq C(\mu^2 + \mu)$, where C is a constant depending only on the equations.

Preuve. Integrating the sum of equations for $c_i = \gamma_a$, we have

$$(5) \quad U_a(x + \gamma_a t, t) = U_a^0(x) + \int_0^t p_a(x + \gamma_a s, s) ds - \int_0^t n_a(x + \gamma_a s, s) ds$$

where

$$(6) \quad p_a(x, t) = \sum' A_{ij}^{k\ell} u_k u_\ell + \sum'' \alpha_i^k u_k,$$

$$(7) \quad n_a(x, t) = \sum_{c_i = \gamma_a} A_{k\ell}^{ij} u_i u_j + \sum_{c_i = \gamma_a} \sum_k \alpha_k^i u_i,$$

where \sum'' [resp. \sum'] is the summation which operates only for $c_k \neq c_i = \gamma_a$ with $\alpha_i^k \neq 0$ [resp. for $c_k \neq c_\ell$ or $c_k = c_\ell$ such that there exist i and j with $A_{ij}^{k\ell} \neq 0$ and $c_k \neq c_i = \gamma_a$]. To prove the global existence of solutions, we knew [18] that $\int_0^\infty \int_{\mathbf{R}} p_a(x, t) dx dt \leq C(\mu^2 + \mu)$.

Theorem 2.— Suppose the conditions (mvQ) and (mvL). Let u_i^0 be positive, summable and bounded Cauchy data. Then the $U_a(x + \gamma_a t, t)$ converge almost everywhere and in L^q with $q \in [1, \infty)$ to a function $\varphi_a(x)$ positive, summable and bounded.

Furthermore, if we suppose the

Condition vd.—

$$i \neq j \Rightarrow c_i \neq c_j,$$

then the $u_i(x + c_i t, t)$ converge almost everywhere and in L^q with $q \in [1, \infty)$ to a function $\varphi_i(x)$.

Preuve. By virtue of Lemma 1, we have

$$(8) \quad U_a(x + \gamma_a t, t) \leq G_a(x) \text{ almost everywhere}$$

where $G_a(x) = U_a^0(x) + \int_0^\infty p_a(x + \gamma_a t, t) dt \geq 0$. Then we have $G_a \in L^1$. We see easily that

$$(9) \quad \begin{aligned} 0 &\leq \int_0^t n_a(x + \gamma_a s, s) ds \\ &\leq U_a^0(x) + \int_0^t p_a(x + \gamma_a s, s) ds \\ &\leq G_a(x). \end{aligned}$$

Since G_a is summable, we have, for almost all $x \in \mathbf{R}$,

$$(10) \quad \int_0^\infty p_a(x + \gamma_a t, t) dt < \infty$$

and

$$(11) \quad \int_0^\infty n_a(x + \gamma_a t, t) dt < \infty.$$

Furthermore we have

$$(12) \quad \int_0^\infty \int_{\mathbf{R}} n_a(x + \gamma_a t, t) dt dx < \infty.$$

We see that

$$(13) \quad \lim_{t \rightarrow +\infty} U_a^0(x) + \int_0^t p_a(x + \gamma_a s, s) ds - \int_0^t n_a(x + \gamma_a s, s) ds,$$

denoted by $\varphi_a(x)$, exists for almost all $x \in \mathbf{R}$. We obtain

$$(14) \quad U_a(\cdot + \gamma_a t, t) \xrightarrow{t \rightarrow +\infty} \varphi_a(\cdot) \text{ almost everywhere.}$$

We see easily that φ_a is positive, summable and bounded. We have

$$(15) \quad \begin{aligned} 0 &\leq U_a^0(x) + \int_0^t p_a(x + \gamma_a s, s) ds - \int_0^t n_a(x + \gamma_a s, s) ds \\ &\leq G_a(x) \in L^1. \end{aligned}$$

By virtue of Lebesgue's theorem, we deduce that

$$(16) \quad U_a^0(\cdot) + \int_0^t p_a(\cdot + \gamma_a s, s) ds - \int_0^t n_a(\cdot + \gamma_a s, s) ds$$

converges to φ_a in $L^1(\mathbf{R})$.

To prove the convergence in L^q with $q \in [1, \infty)$, we have only to use the interpolation between L^1 and L^∞ . Indeed we have

$$(17) \quad \left\| \sum_{i \in E_a} u_i(\cdot + c_i t, t) - \varphi_a(\cdot) \right\|_{L^\infty} \leq C \sup_{\mathbf{R} \times [0, \infty)} \sum_i u_i < \infty.$$

In the above proof, we showed the

Corollary 3.— Suppose the conditions (mvQ) and (mvL). Let u_i^0 be positive, summable and bounded Cauchy data. Then there exists a function G_a positive and summable such that we have

$$(18) \quad U_a(x, t) \leq G_a(x - \gamma_a t)$$

for all $(x, t) \in \mathbf{R} \times [0, \infty)$.

To study better the asymptotic behavior of solutions, we exclude henceforth the case of multiple velocities. The final aim is to show that, for $i \in I_0$, the $u_i(x + c_i t, t)$ tend to $\varphi_i(x)$ uniformly in $x \in \mathbf{R}$. First we have

Proposition 4.— Suppose the conditions (vd), (mvQ) and (mvL). Let u_i^0 be positive, summable and bounded Cauchy data. Then, for all $\varepsilon > 0$, there exists a big T such that we have

$$(19) \quad \int_T^\infty \int_{\mathbf{R}} u_i(x, t) u_j(x, t) dx dt \leq \varepsilon$$

if $i \neq j$,

$$(20) \quad \int_T^\infty \int_{\mathbf{R}} u_i(x, t) dx dt \leq \varepsilon$$

if $i \in I_0$, and

$$(21) \quad \int_{\mathbf{R}} u_i(x, T) dx \leq \varepsilon$$

if $i \in I_0$.

Preuve. By virtue of the remarks concerning the global existence of solutions, we deduce that, for all $\varepsilon > 0$, there exists a big T^0 such that we have

$$(22) \quad \int_{T^0}^\infty \int_{\mathbf{R}} u_i(x, t) u_j(x, t) dx dt \leq \varepsilon$$

if $i \neq j$,

$$(23) \quad \int_{T^0}^\infty \int_{\mathbf{R}} u_i(x, t) dx dt \leq \varepsilon$$

if $i \in I_0$. Since we have

$$(24) \quad \int_{T^0}^{T^0+1} \int_{\mathbf{R}} u_i(x, t) dx dt \leq \varepsilon,$$

there exists a $T \in [T^0, T^0 + 1]$ such that we have

$$(25) \quad \int_{\mathbf{R}} u_i(x, T) dx \leq \varepsilon.$$

We have then the third inequality.

Proposition 5.— Suppose the conditions (vd), (mvQ) and (mvL). Let u_i^0 be positive, summable and bounded Cauchy data. Then, for all $\varepsilon > 0$, there exists a big T such that we have, for c_i, c_k, c_ℓ mutually different,

$$(26) \quad \text{ess sup}_x \int_T^\infty u_k u_\ell(x + c_i t, t) dt \leq \varepsilon$$

and, for $k \in I_0$ such that $i \neq k$,

$$(27) \quad \text{ess sup}_x \int_T^\infty u_k(x + c_i t, t) dt \leq \varepsilon.$$

Preuve. We knew that the $G_i(x) = u_i^0(x, t) + \int_0^\infty p_i(x + c_i t, t) dt$ are positive and summable and that $u_i(x, t) \leq G_i(x - c_i t, t)$. For all $\varepsilon > 0$, there exists a closed interval $K \subseteq \mathbf{R}$ such that $\int_{\mathbf{R} \setminus K} G_i(x) dx < \varepsilon$ for all $i \in I$. We put

$$(28) \quad H_i(x) = \begin{cases} 0, & \text{on } K, \\ G_i(x), & \text{otherwise.} \end{cases}$$

Then we have $\|H_k\|_{L^1} < \varepsilon$. In the outside of the compact set $L = \{(x, t) : x - c_k t, x - c_\ell t \in K\}$, we see that

$$(29) \quad u_k u_\ell(x, t) \leq M (H_k(x - c_k t) + H_\ell(x - c_\ell t)),$$

where we put

$$(30) \quad M = \sup_{(x, t) \in \mathbf{R} \times [0, \infty)} \sum_{i \in I} u_i(x, t) < \infty.$$

There exists a big T such that

$$(31) \quad \mathbf{R} \times [T, \infty) \cap L \neq \emptyset.$$

Then we obtain

$$(32) \quad \begin{aligned} & \int_T^\infty u_k u_\ell(x + c_i t, t) dt \\ & \leq M \int_T^\infty (H_k(x - c_k t) + H_\ell(x - c_\ell t)) dt \\ & \leq 2M\varepsilon. \end{aligned}$$

Therefore we showed the first inequality.

Concerning to the second inequality, we have first, for $t > T$,

$$(33) \quad \begin{aligned} u_k(x + c_i t, t) & \leq u_k(x + c_i t - c_k(t - T), T) \\ & + C \sum_{p \neq q} \int_t^T u_p u_q(x + c_i t - c_k(t - \tau), \tau) d\tau \\ & + \sum_{p \neq k} \alpha_k^p \int_T^t u_p(x + c_i t - c_k(t - \tau), \tau) d\tau. \end{aligned}$$

Then we have

$$(34) \quad \begin{aligned} \int_T^\infty u_k(x + c_i t, t) dt & \leq C \int_{\mathbf{R}} u_k(x, T) dx \\ & + C \sum_{p \neq q} \int_T^\infty \int_{\mathbf{R}} u_p u_q dx dt \\ & + \sum_{p \neq k} \alpha_k^p \int_T^\infty \int_{\mathbf{R}} u_p dx dt. \end{aligned}$$

The two first terms in the right-hand side are less than ε by virtue of the Proposition 4. Concerning to the third term, the summation operates only for $p \neq k$ such that $\alpha_k^p \neq 0$, i.e. for $p \in I_0$. This term is then less also than ε by virtue of the Proposition 4.

Now we state the theorem which describes more precisely the asymptotic behavior of solutions.

Theorem 6.— Suppose the conditions (vd), (mvQ) and (mvL). Let u_i^0 be positive, summable and bounded Cauchy data. Then, for $i \in I_0$, the $u_i(x + c_i t, t)$ tend to 0 uniformly as $t \rightarrow +\infty$. In particular, we have $\varphi_i(x) \equiv 0$.

Preuve. We have

$$(35) \quad \begin{aligned} \frac{d}{dt} u_i(x + c_i t, t) &\leq -a u_i(x + c_i t, t) + C \sum_{p \neq q} u_p u_q(x + c_i t, t) \\ &\quad + C \sum_{p \in I_0} u_p(x + c_i t, t) \end{aligned}$$

with $a > 0$. By virtue of the Proposition 5, for all $\varepsilon > 0$, there exists a big T such that

$$(36) \quad \text{ess sup}_x \int_T^t \sum_{p \neq q} u_p u_q(x + c_i \tau, \tau) d\tau < \varepsilon,$$

$$(37) \quad \text{ess sup}_x \int_T^t \sum_{p \in I_0} u_p(x + c_i \tau, \tau) d\tau < \varepsilon.$$

Integrating the inequality (35), we have, for $t > T$,

$$(38) \quad u_i(x + c_i t, t) \leq M e^{-a(t-T)} + 2\varepsilon$$

with

$$(39) \quad M = \sup_{\mathbf{R} \times [0, \infty)} \sum_{i \in I} u_i(x, t) < \infty.$$

There exists a T^0 such that $M e^{-a(t-T)} < \varepsilon$ for all $t > T^0$. We have therefore, for $t > T^0$, $u_i(x + c_i t, t) < 3\varepsilon$. The function $u_i(x + c_i t, t)$ tends, in L^∞ -norm, to 0 i.e. $\varphi_i(x) \equiv 0$.

Finally we apply the argument due to Shizuta and Kawashima [9], [11] to show the decay in $(1+t)^{-\frac{1}{4}}$ of solutions in case “sufficiently” reflection coefficients α_i^k are non zero, which is incompatible with (mvL) and also necessary for the decay of solutions. First of all, we state some conditions.

Condition \perp .— If $\mu_i \in \mathbf{R}$ verify $\sum_i \mu_i L_i(u) = 0$ for all u , then we have $\sum_i \mu_i Q_i(u) = 0$ for all u .

Condition dsp.— There is no eigenvector λ of $\mathcal{C} = \text{diag}(c_1, \dots, c_N)$ such that λ is in the kernel of \mathcal{L}^t , where $\mathcal{L} = (\alpha_i^j - \delta_{ij} \sum_k \alpha_k^i)_{ij}$ and $N = \#I$.

Remarque : The condition (mvL) is incompatible with the condition (dsp). Indeed, the vector (μ_i) such that $\mu_i = 1$ if $i \in E_0$ and $\mu_i = 0$ otherwise, is an eigenvector of \mathcal{C} and is in the kernel of \mathcal{L}^t .

Corollary 7.— The conditions (\perp) and (dsp) are verified if

1) there is at least two distinct velocities and 0 is a simple eigenvalue of \mathcal{L} .

or

2) (very particular case) there is at least two distinct velocities and all $\alpha_i^k (i \neq k)$ are non zero.

Condition $\mu rL'$.—

$$\alpha_i^k = \alpha_k^i \quad \text{for all } i \text{ and } k$$

(microreversibility of reflection).

Shizuta and Kawashima [9], [11] proved the

Proposition 8.— Suppose the conditions (dsp) and $(\mu rL')$. We put $\mathcal{S} = \mathcal{L} - \mathcal{C} \frac{\partial}{\partial x}$. Let be $u^0 \in H^s \cap L^1(\mathbf{R})$ ($s > \frac{1}{2}$). Then we have

$$(40) \quad \|e^{t\mathcal{S}} u^0\|_{H^s} \leq C(1+t)^{-\frac{1}{4}} \|u^0\|_{H^s \cap L^1}.$$

Furthermore, if u^0 is orthogonal to the kernel of \mathcal{L} , then we have

$$(41) \quad \|e^{t\mathcal{S}} u^0\|_{H^s} \leq C(1+t)^{-\frac{3}{4}} \|u^0\|_{H^s \cap L^1}.$$

We show now the decay of solutions of our nonlinear system.

Theorem 9.— Suppose the conditions (\perp) , (dsp) and $(\mu rL')$. Let u^0 be a positive Cauchy data and in $H^s \cap L^1(\mathbf{R})$ ($s > \frac{1}{2}$). Let $u(t)$ be a solution of

$$(42) \quad \left(\frac{\partial}{\partial t} + \mathcal{C} \frac{\partial}{\partial x} \right) u = Q(u) + L(u),$$

with the Cauchy data u^0 , where $Q(u) = (Q_i(u))_i$, $L(u) = (L_i(u))_i$. If u^0 is sufficiently small in $H^s \cap L^1(\mathbf{R})$, then we have the global existence of solution and the decay of solution in $H^s(\mathbf{R})$:

$$(43) \quad \|u(t)\|_{L^\infty} \leq \|u(t)\|_{H^s} \leq C(1+t)^{-\frac{1}{4}} \|u(t)\|_{H^s \cap L^1}$$

where the constant C depends only on the equations.

Preuve. Owing to the usual argument, it is sufficient to show the estimate (43) up to the time-existence of solution T^* . By virtue of the condition (\perp) , $Q(u)$ is orthogonal to the kernel of \mathcal{L}^t . We have

$$(44) \quad u(t) = e^{t\mathcal{S}} u^0 + \int_0^t e^{(t-\tau)\mathcal{S}} Q(u)(\tau) d\tau.$$

Remarking that $H^s(\mathbf{R})$ forms an algebra for $s > \frac{1}{2}$, we obtain, by the Proposition 8,

$$(45) \quad \begin{aligned} & \|u(t)\|_{H^s} \\ & \leq C(1+t)^{-\frac{1}{4}} (\|u^0\|_{H^s} + \|u^0\|_{L^1}) \\ & + C \int_0^t (1+t-\tau)^{-\frac{3}{4}} (\|Q(u)(\tau)\|_{H^s} + \|Q(u)(\tau)\|_{L^1}) d\tau \\ & \leq CU_0(1+t)^{-\frac{1}{4}} + C \int_0^t (1+t-\tau)^{-\frac{3}{4}} \|u(\tau)\|_{H^s}^2 d\tau, \end{aligned}$$

where we put $U_0 = \|u^0\|_{H^s} + \|u^0\|_{L^1}$. Denoting $U(t) = \sup_{\tau \in [0, t]} (1 + \tau)^{\frac{1}{4}} \|u(\tau)\|_{H^s}$, we have

$$(46) \quad \begin{aligned} U(t) &\leq CU_0 + C(1+t)^{\frac{1}{4}} U(t)^2 \int_0^t (1+t-\tau)^{-\frac{3}{4}} (1+\tau)^{-\frac{1}{2}} d\tau \\ &\leq CU_0 + CU(t)^2. \end{aligned}$$

We remark here that the equation $X = CX_0 + CX^2$ admits two real roots α and β ($\alpha \leq \beta$) for sufficiently small X_0 with $\alpha = O(X_0^2)$. By the continuity of $U(t)$, the value $U(t)$ is included in the interval $[0, \alpha]$, which completes the proof.

At the end, we show that the condition (dsp) is also necessary for the decay of solutions under the condition (\perp).

Theorem 10.— Suppose the condition (\perp). Furthermore we suppose that the condition (dsp) is not verified. Let u_i^0 be positive, summable and bounded Cauchy data. Then the solutions do not tend to 0 except when the data verify a linear relation well precise.

Preuve. We know already [13] that the solutions $u_i(x, t)$ are positive, summable in x and locally bounded in $\mathbf{R} \times \mathbf{R}_+$. By hypothesis, there exists $\mu = (\mu_i) \in \ker \mathcal{L}^t$ and $\gamma \in \mathbf{R}$ such that $\mu_i \neq 0 \Rightarrow c_i = \gamma$. Then we have $\sum_i \mu_i L_i(u) \equiv 0$ and, by the condition (\perp), $\sum_i \mu_i Q_i(u) \equiv 0$. We have then

$$(47) \quad \left(\frac{\partial}{\partial t} + \gamma \frac{\partial}{\partial x} \right) \left(\sum_i \mu_i u_i \right) = 0.$$

Therefore $\sum_i \mu_i u_i$ is a conservative quantity which moves at the velocity γ . Except when $\sum_i \mu_i u_i^0 \equiv 0$, the solutions u_i can not tend to 0.

REFERENCES

1. Bony J.M., *Solutions globales bornées pour les modèles discrets de l'équation de Boltzmann, en dimension 1 d'espace*, Actes des journées E.D.P. à Saint-Jean-de-Monts (1987), École Polytechnique.
2. Bony J.M., *Problème de Cauchy et diffusion à données petites pour les modèles discrets de la cinétique des gaz*, Actes des journées E.D.P. à Saint-Jean-de-Monts (1990), Ecole Polytechnique.
3. Bony J.M., *Existence globale et diffusion en théorie cinétique discrète*, Actes du Symposium en l'honneur de Henri Cabannes, Paris (1990).
4. Bony J.M., *Existence globale à données de Cauchy petites pour les modèles discrets de l'équation de Boltzmann*, Comm. in Partial Differential Equations **16** (1991), 533–545.
5. Cabannes H., *The discrete Boltzmann equation*, University of California at Berkely, 1980.
6. Carleman T., *Problèmes mathématiques dans la théorie cinétique des gaz*, Uppsala (1957), Scientific Publications of Mittag-Leffler Institute.
7. Gatignol R., *Théorie cinétique d'un gaz à répartition discrète de vitesses*, vol. **36**, Springer-Verlag, Heidelberg, 1975.
8. Gatignol R.-Soubbaramayer, *Proceedings of a Symposium Held in Honor of Professor Henri Cabannes at the University Pierre et Marie Curie, Paris, France, on 6 July 1990*, Springer-Verlag, Heidelberg.

9. Kawashima S., *Global Existence and Stability of Solutions for Discrete Velocity Models of the Boltzmann Equation*, Recent Topics In Nonlinear PDE, Hiroshima, Lecture Notes in Num. Appl. Anal. **6** (1983), 59–85.
10. Nishida T., Mimura M., *On the Broadwell's model for a simple discrete velocity gas*, Proc. Japan Acad. **50** (1974), 812–817.
11. Shizuta Y., Kawashima S., *Systems of equations of hyperbolic-parabolic type with applications to the discrete Boltzmann equation*, Hokkaido Mathematical Journal **14** (1985), 249–275.
12. Tartar L., Crandall M.G., *Existence globale pour un système hyperbolique de la théorie des cinétique des gaz*, Ecole Polytechnique, Séminaire Goulaouic-Schwartz n°1 (1975).
13. Yamazaki M., *Existence globale pour les modèles discrets de l'équation de Boltzmann dans un tube mince infini*, C.R.Acad.Sci., Série I, Paris **313** (1991), 29–32.
14. Yamazaki M., *Existence globale pour les modèles discrets de l'équation de Boltzmann dans un tube mince infini en dimension 1 d'espace*, J.Fac.Sci.Univ.Tokyo, Sec. IA **39** (1992), 43–59.
15. Yamazaki M., *Etude de l'équation de Boltzmann pour les modèles discrets dans un tube mince infini*, Transport Theory and Statistical Physics, Proceedings of the Fourth MAFPD **21** (1992), 465–473.
16. Yamazaki M., *On the Discrete Boltzmann Equation with Linear and Nonlinear Terms*, Thesis at University of Tokyo (1993).
17. Yamazaki M., *On the Discrete Boltzmann Equation with Linear and Nonlinear Terms*, J.Fac.Sci.Univ.Tokyo (to appear).
18. Yamazaki M., *Sur les modèles discrets de l'équation de Boltzmann avec termes linéaires et quadratiques*, Thesis of Ecole Polytechnique, Palaiseau, France (1993).